

PREMALATHA

T. SOWMYA

Böcher's theorem in a space of dimension one

Annales mathématiques Blaise Pascal, tome 7, n° 1 (2000), p. 81-86

http://www.numdam.org/item?id=AMBP_2000__7_1_81_0

© Annales mathématiques Blaise Pascal, 2000, tous droits réservés.

L'accès aux archives de la revue « Annales mathématiques Blaise Pascal » (<http://math.univ-bpclermont.fr/ambp/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

BÖCHER'S THEOREM IN A SPACE OF DIMENSION ONE

Premalatha and T. Sowmya

Abstract:

In this paper we express a harmonic function h defined outside a compact set in a B.H. space Ω as an integral with respect to a signed measure in Ω assuming Ω satisfies the axiom of local proportionality. If in particular h is positive and Ω has harmonic dimension one then this expression leads to an analogue of Böcher's theorem in a space of dimension one.

AMS Subject Classification: (1991) 31 D 05.

§1. Introduction

We consider a harmonic function h defined outside a compact set in a B.H. space Ω . This can be written as the difference of two superharmonic functions in Ω where both functions have the same compact support in Ω . If we assume the axiom of local proportionality this leads to an integral representation for h with respect to a signed measure which looks like the Riesz representation. This is of interest because the Riesz representation does not give an integral for a harmonic function as the measure associated with a harmonic function is zero. This theorem gives an analogue of Böcher's theorem in a B.H. space of harmonic dimension one if we assume h is positive.

§2. Preliminaries

Let Ω be a harmonic space satisfying the axioms 1,2,3 of M.Brelot. We assume that constants are harmonic in Ω in which case Ω is referred to as a B.H.space. Ω is called a B.P. or B.S. space according as there exists a positive potential or not in Ω . For a nonlocally polar outer regular compact set $k \subset \Omega$ and a continuous function f on ∂k , as in [1], the notation $B_k f$ stands for the Dirichlet solution in $\Omega - k$ with values f on ∂k and 0 at the point at infinity.

In the case of a B.S. space Ω , we fix an outer regular compact set K and a regular domain ω , $K \subset \omega$ with respect to which flux is defined (for definition see [1]). We also fix a harmonic function $H > 0$ in $\Omega - K$ tending to 0 on ∂K with flux at infinity one.

We recall the definition of a B.H. potential in a B.S. space Ω : Let $\{\Omega_i\}$ be a fixed regular exhaustion of Ω . Fix an ultrafilter e finer than the filter of sections of $\{\Omega_i\}$. Let $\mathcal{D}(u)$ be the limit of $\overline{M}_u^{\Omega_i}$ according to the ultrafilter e . An admissible superharmonic function u in a B.S. space Ω with flux at infinity α is said to be a B.H. - potential if $\mathcal{D}(u - \alpha H) = 0$.

It can be easily seen that a superharmonic function u with compact support in a B.P. (respectively B.S.) space can be written uniquely as the sum of a potential (respectively B.H. potential) and a harmonic function.

Let Ω be a B.H. space satisfying the axiom of local proportionality.

Case (i). Let Ω be a B.P. space. If δ is a regular domain and z a fixed point in δ , then for any y there exists a unique potential $q_y(x)$ with support y such that $\int q_y(x) d\rho_z^\delta(x) = 1$ where $d\rho_z^\delta$ is the harmonic measure of δ with respect to z .

If u is a potential with compact support A then there exists a unique Radon measure $\mu \geq 0$ supported by A such that $u(x) = \int q_y(x) d\mu(y)$; and conversely if $\mu \geq 0$ is a Radon measure with compact support then $\int q_y(x) d\mu(y)$ is a potential.

Case (ii): Let Ω be a B.S. space. In this case, for any y , there exists a unique B.H. potential $q_y(x)$ with support y and flux q_y at infinity -1 . Then if $u(x)$ is a B.H. potential with compact support A , there exists a unique Radon measure $\mu \geq 0$ supported by A such that $u(x) = \int q_y(x) d\mu(y)$; and conversely, if $\mu \geq 0$ is a Radon measure with compact support, then $u(x) = \int q_y(x) d\mu(y)$ is a B.H. potential with flux u at infinity $= - \int d\mu$.

§ 3. BÖCHER'S THEOREM IN A SPACE OF DIMENSION ONE

Theorem 1.

Let h be a harmonic function defined outside a compact set X in a B.H. space Ω and ω_0 be any regular domain such that $X \subset \omega_0$. Assume that Ω has a countable base and satisfies the axiom of local proportionality. Then

there exists a signed measure μ with support contained in $\partial\omega_0$ and a uniquely determined harmonic function u in Ω such that $h(x) = \int q_y(x)d\mu(y) + u(x)$ in $\Omega \sim \bar{\omega}_0$.

Here $q_y(x)$ is the potential (respectively B.H. potential) that we fix in Ω as explained in §2, if Ω is a B.P. (respectively B.S.) space. Moreover if the harmonic dimension at infinity of Ω is 1, u is a constant if and only if h is bounded on one side near the point at infinity A .

Proof.

Let $x_0 \in X$ and s_{x_0} be a superharmonic function in Ω with point support x_0 .

Choose an outer regular compact set K_1 such that $X \subset K_1^0 \subset K_1 \subset \omega_0$.

Without loss of generality we can assume that h is harmonic in $\omega_0 \sim K_1$ and continuous in $\bar{\omega}_0 \sim \bar{K}_1$. For a continuous function f on $\partial\omega_0$ let $Df = H_f^{\omega_0}$ denote the Dirichlet solution in ω_0 with boundary value f .

Since $Ds_{x_0} < s_{x_0}$ in ω_0 we have $\inf_{\partial K_1} (s_{x_0} - Ds_{x_0}) > 0$.

Choose $\alpha > 0$ such that

$$\alpha(s_{x_0} - Ds_{x_0}) > Dh - h \text{ on } \partial K_1.$$

Then $h + \alpha s_{x_0} > D(h + \alpha s_{x_0})$ on ∂K_1 .

Since $h + \alpha s_{x_0} = D(h + \alpha s_{x_0})$ on $\partial\omega_0$, by minimum principle of harmonic functions we get

$$h + \alpha s_{x_0} > D(h + \alpha s_{x_0}) \text{ in } \omega_0 \sim K_1.$$

Define
$$h_1 = \begin{cases} h + \alpha s_{x_0} & \text{in } \Omega \sim \omega_0 \\ D(h + \alpha s_{x_0}) & \text{in } \omega_0 \end{cases}$$

and
$$h_2 = \begin{cases} \alpha s_{x_0} & \text{on } \Omega \sim \omega_0 \\ D(\alpha s_{x_0}) & \text{on } \omega_0. \end{cases}$$

Then h_1 and h_2 are finite, continuous, superharmonic functions in Ω with compact support in $\partial\omega_0$ such that

$$h = h_1 - h_2 \text{ on } \Omega \sim \bar{\omega}_0.$$

Now, $h_i = p_i + u_i$ $i = 1, 2$ where p_i is a potential (respectively B.H. potential) with support in $\partial\omega_0$ if Ω is a B.P. (respectively B.S.) space and u_i is harmonic in Ω .

Hence $h = p_1 - p_2 + u$ where $u = u_1 - u_2$ is harmonic in Ω . But $p_i(x) = \int q_y(x) d\mu_i(y)$, $i = 1, 2$ where $\mu_i, i = 1, 2$ is a Radon measure with support contained in $\partial\omega_0$.

Hence $h(x) = \int q_y(x) d\mu(y) + u(x)$ where $\mu = \mu_1 - \mu_2$ is a signed measure with support contained in $\partial\omega_0$.

We shall complete the proof by considering the two cases of a B.P. space and a B.S. space separately.

Case (i). Let Ω be a B.P. space.

Suppose $h(x) = \int q_y(x) d\mu'(y) + u'(x)$ where μ' is also a signed measure with support contained in $\partial\omega_0$ and u' is harmonic in Ω .

Then h can be written as

$$h = p_1 - p_2 + u = q_1 - q_2 + u' \text{ in } \Omega \sim \bar{\omega}_0$$

where $q_i, i = 1, 2$ are potentials in Ω with compact support.

$$\begin{aligned} \text{Then } \mathcal{D}(p_1) &= \mathcal{D}(p_2) = \mathcal{D}(q_1) = \mathcal{D}(q_2) = 0 \text{ gives} \\ \mathcal{D}(u) &= u = u' = \mathcal{D}(u'). \end{aligned}$$

Thus u is uniquely determined in Ω .

Now $h = p_1 - p_2 + u$ on $\Omega \sim \bar{\omega}_0$.

Since p_1 and p_2 are potentials with compact support, they are bounded outside a compact set in Ω .

Hence if h is bounded on one side near \mathcal{A} so is u .

Therefore if Ω is of harmonic dimension one, we see that u reduces to a constant [2].

If u is a constant then clearly h is bounded on one side near \mathcal{A} .

Case (ii): Let Ω be a B.S. space.

Let flux $p_1 = \alpha_1$ and flux $p_2 = \alpha_2$.

Then $h - (\alpha_1 - \alpha_2)H = (p_1 - \alpha_1 H) - (p_2 - \alpha_2 H) + u$

gives $\mathcal{D}(h - (\alpha_1 - \alpha_2)H) = u$ by definition of a B.H. potential.

Since $\alpha_1 - \alpha_2 = \text{flux } h$, we see that given h, u is uniquely determined in Ω .

Now since $D(p_i - \alpha_i H) = 0$ we get $p_i - \alpha_i H$, $i = 1, 2$ are bounded outside a compact set.

Hence $h = u + (\alpha_1 - \alpha_2)H +$ a bounded harmonic function outside a compact set.

If h is bounded on one side near \mathcal{A} , then $u + (\alpha_1 - \alpha_2)H$ is bounded on one side near \mathcal{A} .

If Ω has harmonic dimension one this implies that u is a constant [2].

If u is a constant, h is obviously bounded on one side near \mathcal{A} .

This completes the proof of the theorem.

Now if we take the function h in the above theorem to be ≥ 0 we can deduce the analogue of the inverted version of Böcher's theorem, which may be stated as follows, in a space of harmonic dimension one.

Böcher's theorem: (Inverted version). Let u be positive and harmonic in $\mathbb{R}^n - \bar{B}$, $n \geq 2$ where B is the unit ball about the origin. Then

$$u(x) = \begin{cases} \alpha \log|x| + b(x) & \text{if } n = 2 \\ \alpha + b(x) & \text{if } n \geq 3 \end{cases}$$

where $b(x)$ is a bounded harmonic function in $\mathbb{R}^n - \bar{B}$ and $\alpha \geq 0$ is a constant. If $n \geq 3$, $b(x)$ is actually bounded by a bounded potential.

This can be proved by applying the Kelvin's transform to the standard form of Böcher's theorem [3].

Theorem 2.

Let Ω be a B.H. space of harmonic dimension one and h be a positive harmonic function defined outside a compact set X . If Ω is a B.P. space then $h = \alpha + b$ where α is a constant and b is a harmonic function bounded by a bounded potential outside a compact set.

If Ω is B.S., then $h = \alpha H + b$ outside a compact set where α is a constant and b is a bounded harmonic function outside a compact set.

Proof.

Case (i). Let Ω be a B.P. space.

Take ω_0, p_1, p_2 as in Theorem 1.

Since $h \geq 0$, u is a constant say α .

Let K' be an outer regular compact set such that $(K')^0 \supset \partial\omega_0$.

Then for $i = 1, 2$, $v_i = \begin{cases} 0 & \text{on } K' \\ p_i - B_{K'}p_i & \text{on } \Omega \sim K' \end{cases}$
 is a subharmonic function on Ω such that $0 \leq v_i \leq p_i$.

Since p_i is a potential this implies that $v_i \equiv 0$ or $p_i = B_{K'}p_i$ outside the compact set K' .

If $p_i \leq \lambda$ on $\partial K'$, then $B_{K'}p_i \leq \lambda B_{K'}1$.

Hence $h = p_1 - p_2 + \alpha = \alpha + b$ where $b = p_1 - p_2$ is such that $|b| \leq 2\lambda B_{K'}1$, a bounded potential outside a compact set.

Case (ii). Let Ω be a B.S. space.

Then as in the proof of the above theorem since u is a constant we get

$$\begin{aligned} h &= (\alpha_1 - \alpha_2)H + \text{a bounded harmonic function} \\ &= \alpha H + b \quad \text{outside a compact set.} \end{aligned}$$

References

1. V. Anandam, Espaces Harmoniques Sans Potentiel Positif, Ann. Inst. Fourier, Grenoble, 22, 4 (1972) 97-160.
2. V. Anandam, Potentials in a B.S. Harmonic Space, Bull. Math. Roumanie, 18 3-4 (1974), 233 - 248.
3. S. Axler, T. Bourdon and W. Ramey, Harmonic Function Theory, GTM, Vol. 137, Springer-Verlag, Germany, 993.

Premalatha

*Ramanujan Institute for Advanced Study in Mathematics
University of Madras, CHENNAI 600 005, India.*

T. Sowmya

*Ramanujan Institute for Advanced Study in Mathematics
University of Madras, CHENNAI 600 005, India.*